


## Trigonometry Review

Consider a right angle triangle



Let  $\angle BAC = \theta$ ,

Side  $AB = c$  and side  $BC = a$ .

We can calculate Side  $AC = b$ , by Pythagorean Theorem

$$a^2 + b^2 = c^2$$

**We can define 6 trigonometric ratios :**

$$\sin \theta = \frac{\text{Opposite Side}}{\text{Hypotenuse}} = \frac{a}{c}$$

$$\csc \theta = \frac{\text{Hypotenuse}}{\text{Opposite Side}} = \frac{c}{a}$$

$$\cos \theta = \frac{\text{Adjacent Side}}{\text{Hypotenuse}} = \frac{b}{c}$$

$$\sec \theta = \frac{\text{Hypotenuse}}{\text{Adjacent Side}} = \frac{c}{b}$$

$$\tan \theta = \frac{\text{Opposite Side}}{\text{Adjacent Side}} = \frac{a}{b}$$

$$\cot \theta = \frac{\text{Adjacent Side}}{\text{Opposite Side}} = \frac{b}{a}$$

**From these definitions we can derive:**

$$\sin \theta = \frac{1}{\csc \theta} \qquad \csc \theta = \frac{1}{\sin \theta}$$

$$\cos \theta = \frac{1}{\sec \theta} \qquad \sec \theta = \frac{1}{\cos \theta}$$

$$\tan \theta = \frac{1}{\cot \theta} \qquad \cot \theta = \frac{1}{\tan \theta}$$

Also,

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \qquad \cot \theta = \frac{\cos \theta}{\sin \theta}$$


**For Example**

We have previously defined

$$\sin \theta = \frac{a}{c} \quad , \quad \frac{1}{\sin \theta} = \frac{c}{a} = \csc \theta$$

This can be done for the other trigonometric functions as well.


## Signs of function values




A trick to remember this would be to remember the phrase “All Silver Tea Cups” or “A Smart Trig Class,” starting in the first quadrant.

## To find function values of special angles

Consider an equilateral triangle, length of each side being 2 units (see figure). Let us bisect this triangle as shown. We will end up with two 30°-60° triangles.



Now let us consider one of the bisected triangles



By Pythagorean theorem,  
 $1^2 + (BC)^2 = 2^2$ ,  $(BC)^2 = 4 - 1 = 3$   
 Therefore,  $BC = \sqrt{3}$

We can now find values for 30°.

$$\sin 30^\circ = \frac{\text{Opposite Side}}{\text{Hypotenuse}} = \frac{1}{2} \qquad \cos 30^\circ = \frac{\text{Adjacent Side}}{\text{Hypotenuse}} = \frac{\sqrt{3}}{2}$$


$$\tan 30^\circ = \frac{\text{Opposite Side}}{\text{Adjacent Side}} = \frac{1}{\sqrt{3}} = \frac{\sqrt{3}}{3}$$

Similarly we can find values for 60°.

$$\sin 60^\circ = \frac{\text{Opposite Side}}{\text{Hypotenuse}} = \frac{\sqrt{3}}{2} \qquad \cos 60^\circ = \frac{\text{Adjacent Side}}{\text{Hypotenuse}} = \frac{1}{2}$$

$$\tan 60^\circ = \frac{\text{OppositeSide}}{\text{AdjacentSide}} = \sqrt{3}$$

To find values for  $45^\circ$ , we consider a  $45^\circ$ - $45^\circ$  triangle



By Pythagorean Theorem

$$(AB)^2 = 1^2 + 1^2 = 2$$

$$AB = \sqrt{2}$$

$$\sin 45^\circ = \frac{\text{OppositeSide}}{\text{Hypotenuse}} = \frac{1}{\sqrt{2}} = \frac{\sqrt{2}}{2}$$


$$\cos 45^\circ = \frac{\text{AdjacentSide}}{\text{Hypotenuse}} = \frac{1}{\sqrt{2}} = \frac{\sqrt{2}}{2}$$

$$\tan 45^\circ = \frac{\text{OppositeSide}}{\text{AdjacentSide}} = 1$$

**Table for the most commonly used angles**

Angle $\theta$	Sine	Cosine	Tangent
$0^\circ$	0	1	0
$30^\circ$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$
$45^\circ$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1
$60^\circ$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$
$90^\circ$	1	0	Undefined
$180^\circ$	0	-1	0

**Let us now derive some trigonometric identities:**



By the Pythagorean Theorem,

$$a^2 + b^2 = c^2 \quad (1.1)$$

From earlier definitions we have

$$\sin \theta = \frac{a}{c}, \quad \cos \theta = \frac{b}{c}, \quad \tan \theta = \frac{a}{b}, \quad \csc \theta = \frac{c}{a}, \quad \sec \theta = \frac{c}{b}, \quad \cot \theta = \frac{b}{a}$$

**The three trigonometric identities:**

Let us divide equation 1.1 by  $c^2$ ,

We get

$$\frac{a^2}{c^2} + \frac{b^2}{c^2} = \frac{c^2}{c^2}$$

$$\frac{a^2}{c^2} + \frac{b^2}{c^2} = 1$$

$$(\sin \theta)^2 + (\cos \theta)^2 = 1 \quad \text{or} \quad \sin^2 \theta + \cos^2 \theta = 1$$

If we divide equation 1.1 by  $a^2$  we get,

$$1 + \frac{b^2}{a^2} = \frac{c^2}{a^2}$$

$$1 + (\cot \theta)^2 = (\csc \theta)^2 \quad \text{or} \quad 1 + \cot^2 \theta = \csc^2 \theta$$

Finally we divide equation 1.1 by  $b^2$  we get,

$$\frac{a^2}{b^2} + 1 = \frac{c^2}{b^2}$$

$$(\tan \theta)^2 + 1 = (\sec \theta)^2 \quad \text{or} \quad \tan^2 \theta + 1 = \sec^2 \theta$$